

# AN OPERATOR EQUALITY INVOLVING A CONTINUOUS FIELD OF OPERATORS AND ITS NORM INEQUALITIES

MOHAMMAD SAL MOSLEHIAN AND FUZHEN ZHANG

**ABSTRACT.** Let  $\mathfrak{A}$  be a  $C^*$ -algebra,  $T$  be a locally compact Hausdorff space equipped with a probability measure  $P$  and let  $(A_t)_{t \in T}$  be a continuous field of operators in  $\mathfrak{A}$  such that the function  $t \mapsto A_t$  is norm continuous on  $T$  and the function  $t \mapsto \|A_t\|$  is integrable. Then the following equality including Bouchner integrals holds

$$\int_T \left| A_t - \int_T A_s dP \right|^2 dP = \int_T |A_t|^2 dP - \left| \int_T A_t dP \right|^2. \quad (0.1)$$

This equality is related both to the notion of variance in statistics and to a characterization of inner product spaces. With this operator equality, we present some uniform norm and Schatten  $p$ -norm inequalities.

## 1. INTRODUCTION AND PRELIMINARIES

Many interesting characterizations of inner product spaces have been introduced (see, e.g., [1]). It is shown by Th. M. Rassias [8] that a normed space  $X$  (with norm  $\|\cdot\|$ ) is an inner product space if and only if for any finite set of vectors  $x_1, \dots, x_n \in X$ ,

$$\sum_{i=1}^n \left\| x_i - \frac{1}{n} \sum_{j=1}^n x_j \right\|^2 = \sum_{i=1}^n \|x_i\|^2 - n \left\| \frac{1}{n} \sum_{j=1}^n x_j \right\|^2. \quad (1.1)$$

This equality is of fundamental importance in the study of normed spaces and inner product spaces since it reveals a basic relation between the two sorts of spaces. From statistical point of view, let  $(X, \mu)$  be a probability measure space and  $f$  be a random variable, i.e.,  $f$  is an element of  $L^2(X, \mu)$ . With the variance of  $f$  defined by

$$\text{Var}(f) = E(|f - E(f)|^2),$$

---

2000 *Mathematics Subject Classification.* Primary 47A62; secondary 46C15, 47A30, 15A24.

*Key words and phrases.* Bounded linear operator; Characterization of inner product space; Hilbert space;  $Q$ -norm; Norm inequality; Schatten  $p$ -norm; continuous field of operators, Bouchner integral.

where  $E(f) = \int_X f d\mu$  denotes the expectation of  $f$ , (1.1) resembles (or vice versa) the well-known equality

$$\text{Var}(f) = E(|f|^2) - |E(f)|^2.$$

Let  $\mathbb{B}(\mathcal{H})$  be the algebra of all bounded linear operators on a separable complex Hilbert space  $\mathcal{H}$  endowed with inner product  $\langle \cdot, \cdot \rangle$  and the operator norm  $\|\cdot\|_\infty$ . We denote the absolute value of  $A \in \mathbb{B}(\mathcal{H})$  by  $|A| = (A^*A)^{1/2}$ . For  $x, y \in \mathcal{H}$ , the rank one operator  $x \otimes y$  is defined on  $\mathcal{H}$  by  $(x \otimes y)(z) = \langle z, y \rangle x$ .

Let  $\mathfrak{A}$  be a  $C^*$ -algebra and let  $T$  be a locally compact Hausdorff space. A field  $(A_t)_{t \in T}$  of operators in  $\mathfrak{A}$  is called a continuous field of operators if the function  $t \mapsto A_t$  is norm continuous on  $T$ . If  $\mu(t)$  is a Radon measure on  $T$  and the function  $t \mapsto \|A_t\|$  is integrable, one can form the Bochner integral  $\int_T A_t d\mu(t)$ , which is the unique element in  $\mathfrak{A}$  such that

$$\varphi \left( \int_T A_t d\mu(t) \right) = \int_T \varphi(A_t) d\mu(t)$$

for every linear functional  $\varphi$  in the norm dual  $\mathfrak{A}^*$  of  $\mathfrak{A}$ ; cf. [6, Section 4.1].

Let  $A \in \mathbb{B}(\mathcal{H})$  be a compact operator and let  $0 < p < \infty$ . The Schatten  $p$ -norm ( $p$ -quasi-norm) for  $1 \leq p < \infty$  ( $0 < p < 1$ ) is defined by  $\|A\|_p = (\text{tr}|A|^p)^{1/p}$ , where  $\text{tr}$  is the usual trace functional. Clearly, for  $p, q > 0$ ,

$$\|A\|_p^q = \| |A|^2 \|_{p/2}^{q/2}.$$

For  $p > 0$ , the Schatten  $p$ -class, denoted by  $C_p$ , is defined to be the two-sided ideal in  $\mathbb{B}(\mathcal{H})$  of those compact operators  $A$  for which  $\|A\|_p$  is finite. In particular,  $C_1$  and  $C_2$  are the trace class and the Hilbert-Schmidt class, respectively. For more information on the theory of the Schatten  $p$ -classes, the reader is referred to [2, 9].

Since  $C_2$  is a Hilbert space under the inner product  $\langle A, B \rangle = \text{tr}(B^*A)$ , it follows from (1.1) that if  $A_1, \dots, A_n \in C_2$ , then

$$\sum_{i=1}^n \left\| A_i - \frac{1}{n} \sum_{j=1}^n A_j \right\|_2^2 = \sum_{i=1}^n \|A_i\|_2^2 - n \left\| \frac{1}{n} \sum_{j=1}^n A_j \right\|_2^2. \quad (1.2)$$

In this paper, we establish a general operator version of equality (1.1), from which we deduce an extension of (1.1). We present some inequalities concerning various norms such as Schatten  $p$ -norms that form natural generalizations (in inequalities) of the identity (1.2) (see, e.g., [7]). It seems that the inequalities related to Schatten  $p$ -norms are useful in operator theory and mathematical physics and are interesting in their own right.

## 2. MAIN RESULTS

We begin by establishing an operator version of equality (1.1) involving continuous fields of operators and integral means of operators.

**Theorem 2.1.** *Let  $\mathfrak{A}$  be a  $C^*$ -algebra,  $T$  be a locally compact Hausdorff space equipped with a probability measure  $P$  and let  $(A_t)_{t \in T}$  be a continuous field of operators in  $\mathfrak{A}$  such that the function  $t \mapsto A_t$  is norm continuous on  $T$  and the function  $t \mapsto \|A_t\|$  is integrable. Then*

$$\int_T \left| A_t - \int_T A_s dP(s) \right|^2 dP(t) = \int_T |A_t|^2 dP(t) - \left| \int_T A_t dP(t) \right|^2 \quad (2.1)$$

*Proof.*

$$\begin{aligned} & \int_T \left| A_t - \int_T A_s dP(s) \right|^2 dP(t) \\ &= \int_T \left( A_t - \int_T A_s dP(s) \right)^* \left( A_t - \int_T A_r dP(r) \right) dP(t) \\ &= \int_T |A_t|^2 dP(t) - \int_T \left( A_t^* \int_T A_r dP(r) \right) dP(t) \\ &\quad - \int_T \left( \left( \int_T A_s dP(s) \right)^* A_t \right) dP(t) \\ &\quad + \left( \int_T A_s dP(s) \right)^* \int_T A_r dP(r) \int_T dP(t) \\ &= \int_T |A_t|^2 dP(t) - \left( \int_T A_s dP(s) \right)^* \int_T A_r dP(r) \\ &\quad \left( \text{by } \int_T dP(t) = 1 \text{ and } \int_T A_t^* dP(t) = \left( \int_T A_t dP(t) \right)^* \right) \\ &= \int_T |A_t|^2 dP(t) - \left| \int_T A_t dP(t) \right|^2. \end{aligned}$$

□

**Corollary 2.2.** *Let  $A_1, \dots, A_n \in \mathbb{B}(\mathcal{H})$ . Then*

(i) *for any set of nonnegative numbers  $t_1, \dots, t_n$  with  $\sum_{i=1}^n t_i = 1$ ,*

$$\sum_{i=1}^n t_i \left| A_i - \sum_{j=1}^n t_j A_j \right|^2 = \sum_{i=1}^n t_i |A_i|^2 - \left| \sum_{j=1}^n t_j A_j \right|^2; \quad (2.2)$$

(ii)

$$\sum_{i=1}^n \left| A_i - \frac{1}{n} \sum_{j=1}^n A_j \right|^2 = \sum_{i=1}^n |A_i|^2 - n \left| \frac{1}{n} \sum_{j=1}^n A_j \right|^2. \quad (2.3)$$

*Proof.* (i) Take  $T = \{1, \dots, n\}$  and  $P(\{i\}) = t_i$  in Theorem 2.1.

(ii) Put  $t_i = 1/n$  in (i).  $\square$

As an immediate consequence of (2.2), we get the following known AM-QM operator inequality; cf. [10, Theorem 7].

**Corollary 2.3.** *Let  $A_1, \dots, A_n \in \mathbb{B}(\mathcal{H})$ . Then for any set of nonnegative numbers  $t_1, \dots, t_n$  with  $\sum_{i=1}^n t_i = 1$ ,*

$$\left| \sum_{i=1}^n t_i A_i \right|^2 \leq \sum_{i=1}^n t_i |A_i|^2.$$

The following result compares the mean of the squares of the operators to the square of the mean, and gives some bounds of their difference.

**Corollary 2.4.** *Let  $A_1, \dots, A_n \in \mathbb{B}(\mathcal{H})$  be positive operators such that  $0 \leq m_i I \leq A_i \leq M_i I$  for some nonnegative scalars  $m_i$  and  $M_i$  and all  $1 \leq i \leq n$ , and let  $t_1, \dots, t_n$  be nonnegative numbers such that  $\sum_{i=1}^n t_i = 1$ . Then*

$$\sum_{i=1}^n t_i \beta_i^2 I \leq \sum_{i=1}^n t_i A_i^2 - \left( \sum_{i=1}^n t_i A_i \right)^2 \leq \sum_{i=1}^n t_i \alpha_i^2 I,$$

where

$$\alpha_i = \max \left\{ \left| M_i - \sum_{i=1}^n t_i m_i \right|, \left| m_i - \sum_{i=1}^n t_i M_i \right| \right\}$$

and

$$\beta_i = \min \left\{ \left| M_i - \sum_{i=1}^n t_i m_i \right|, \left| m_i - \sum_{i=1}^n t_i M_i \right| \right\}.$$

*Proof.* It is sufficient to notice that for each  $A_i$ , by the functional calculus,

$$\begin{aligned} \min \left\{ \left| M_i - \sum_{i=1}^n t_i m_i \right|^2, \left| m_i - \sum_{i=1}^n t_i M_i \right|^2 \right\} &\leq \left| A_i - \sum_{j=1}^n t_j A_j \right|^2 \\ &\leq \max \left\{ \left| M_i - \sum_{i=1}^n t_i m_i \right|^2, \left| m_i - \sum_{i=1}^n t_i M_i \right|^2 \right\} \end{aligned}$$

and use (2.2).  $\square$

By a  $Q$ -norm on a subspace  $\mathcal{D}$  of  $\mathbb{B}(\mathcal{H})$  we mean a norm  $\|\cdot\|_Q$  for which there exists a norm  $\|\cdot\|_{\widehat{Q}}$  defined on a subspace  $\widehat{\mathcal{D}}$  such that  $\|A\|_Q^2 = \|A^*A\|_{\widehat{Q}}$  ( $A \in \mathcal{D}$ ) (we implicitly assume that  $A^*A \in \widehat{\mathcal{D}}$  if and only if  $A \in \mathcal{D}$ ). The operator norm  $\|\cdot\|_\infty$  and  $\|\cdot\|_p$  (for  $2 \leq p < \infty$ ) are examples of  $Q$ -norms on  $\mathbb{B}(\mathcal{H})$ ; see, e.g., [2, p. 89] or [4].

**Corollary 2.5.** *Let  $\|\cdot\|_Q$  be a  $Q$ -norm on a subspace  $\mathcal{D}$  of  $\mathbb{B}(\mathcal{H})$ , and let  $t_1, \dots, t_n$  be nonnegative numbers such that  $\sum_{i=1}^n t_i = 1$ . Then for any  $A_1, \dots, A_n \in \mathcal{D}$  with  $A_i^*A_j = 0$  for  $1 \leq i \neq j \leq n$ ,*

$$\left\| \sum_{i=1}^n \sqrt{t_i} A_i \right\|_Q^2 \leq \sum_{i=1}^n t_i \left\| A_i - \sum_{j=1}^n t_j A_j \right\|_Q^2 + \left\| \sum_{j=1}^n t_j A_j \right\|_Q^2.$$

*Proof.*

$$\begin{aligned} \left\| \sum_{i=1}^n \sqrt{t_i} A_i \right\|_Q^2 &= \left\| \left| \sum_{i=1}^n \sqrt{t_i} A_i \right|^2 \right\|_{\widehat{Q}} \\ &= \left\| \sum_{i=1}^n t_i |A_i|^2 \right\|_{\widehat{Q}} \\ &\leq \sum_{i=1}^n \left\| t_i \left| A_i - \sum_{j=1}^n t_j A_j \right|^2 \right\|_{\widehat{Q}} + \left\| \left| \sum_{j=1}^n t_j A_j \right|^2 \right\|_{\widehat{Q}} \\ &= \sum_{i=1}^n t_i \left\| A_i - \sum_{j=1}^n t_j A_j \right\|_Q^2 + \left\| \sum_{j=1}^n t_j A_j \right\|_Q^2. \end{aligned}$$

□

*Remark 2.6.* The operators  $A_i$  acting on a Hilbert space having the orthogonal property  $A_i^*A_j = 0$  for  $1 \leq i \neq j \leq n$  are not uncommon. For instance, let  $(e_i)$  be an orthogonal family (not containing zero) in  $\mathcal{H}$  and define the operators  $A_i : \mathcal{H} \rightarrow \mathcal{H}$  by  $A_i = \frac{e_i \otimes e_i}{\|e_i\|}$ ,  $1 \leq i \leq n$ . Then  $A_i$ 's are positive operators in  $\mathbb{B}(\mathcal{H})$  with  $\|A_i\|_\infty = \|e_i\|$  for all  $i$  and  $\|A_i A_j\|_\infty = |\langle e_i, e_j \rangle|$  for all  $1 \leq i, j \leq n$  (for details see [5]).

In the setting of Hilbert spaces, the known equality (1.1) can be proved directly. In what follows we show that an extension of it can also be obtained from equality (2.2).

**Corollary 2.7.** *Let  $x_1, \dots, x_n \in \mathcal{H}$ . Then*

$$\sum_{i=1}^n t_i \left\| x_i - \sum_{j=1}^n t_j x_j \right\|^2 = \sum_{i=1}^n t_i \|x_i\|^2 - \left\| \sum_{j=1}^n t_j x_j \right\|^2.$$

*Proof.* Let  $e$  be a non-zero vector of  $\mathcal{H}$  and set  $A_i = x_i \otimes e$ . It follows from the elementary properties of rank one operators and equality (2.2) that

$$\begin{aligned}
\sum_{i=1}^n t_i \left\| x_i - \sum_{j=1}^n t_j x_j \right\|^2 e \otimes e &= \sum_{i=1}^n t_i \left| \left( x_i - \sum_{j=1}^n t_j x_j \right) \otimes e \right|^2 \\
&= \sum_{i=1}^n t_i \left| A_i - \sum_{j=1}^n t_j A_j \right|^2 \\
&= \sum_{i=1}^n t_i |A_i|^2 - \left| \sum_{j=1}^n t_j A_j \right|^2 \\
&= \sum_{i=1}^n t_i |x_i \otimes e|^2 - \left| \left( \sum_{j=1}^n t_j x_j \right) \otimes e \right|^2 \\
&= \left( \sum_{i=1}^n t_i \|x_i\|^2 - \left\| \sum_{j=1}^n t_j x_j \right\|^2 \right) e \otimes e,
\end{aligned}$$

from which we conclude the result.  $\square$

From now on we restrict ourselves to (2.3).

A special case (for  $X = \mathbb{C}$ ) of equality (1.1) is

$$\sum_{i=1}^n \left| z_i - \frac{1}{n} \sum_{j=1}^n z_j \right|^2 = \sum_{i=1}^n |z_i|^2 - n \left| \frac{1}{n} \sum_{j=1}^n z_j \right|^2 \quad (z_1, \dots, z_n \in \mathbb{C}).$$

This equality is in turn a special case (when  $A = \text{diag}(z_1, \dots, z_n)$  and  $\text{tr}([a_{ij}]) := \sum_{i=1}^n a_{ii}$ ) of the next equality concerning the usual normalized trace functional.

**Theorem 2.8.** *Let  $\widehat{\text{tr}}(A) = \text{tr}(A)/\text{tr}(I)$  be the normalized trace on  $M_n(\mathbb{C})$ . Then*

$$\|A - \widehat{\text{tr}}(A)\|_2^2 = \|A\|_2^2 - \text{tr}(I) |\widehat{\text{tr}}(A)|^2.$$

*Proof.*

$$\begin{aligned}
\|A - \widehat{\text{tr}}(A)\|_2^2 &= \text{tr} \left| A - \frac{\text{tr}(A)}{\text{tr}(I)} I \right|^2 \\
&= \text{tr} \left( \left( A^* - \frac{\overline{\text{tr}(A)}}{\text{tr}(I)} I \right) \left( A - \frac{\text{tr}(A)}{\text{tr}(I)} I \right) \right) \\
&= \text{tr}(A^* A) - \frac{\text{tr}(A^*) \text{tr}(A)}{\text{tr}(I)} - \frac{\overline{\text{tr}(A)} \text{tr}(A)}{\text{tr}(I)} + \frac{|\text{tr}(A)|^2}{\text{tr}(I)^2} \text{tr}(I) \\
&= \text{tr}|A|^2 - \text{tr}(I) \left| \frac{\text{tr}(A)}{\text{tr}(I)} \right|^2 \quad (\text{by } \text{tr}(A^*) = \overline{\text{tr}(A)}) \\
&= \|A\|_2^2 - \text{tr}(I) |\widehat{\text{tr}}(A)|^2.
\end{aligned}$$

□

We need the following lemma which can be deduced from [4, Lemma 4] and [9, p. 20] (see also [7, Lemma 2.1]).

**Lemma 2.9.** *Let  $A_1, \dots, A_n$  be positive operators in  $C_p$  for some  $p > 0$ .*

(i) *If  $0 < p < 1$ , then*

$$n^{p-1} \sum_{i=1}^n \|A_i\|_p^p \leq \left\| \sum_{i=1}^n A_i \right\|_p^p \leq \sum_{i=1}^n \|A_i\|_p^p.$$

(ii) *If  $1 \leq p < \infty$ , then*

$$\sum_{i=1}^n \|A_i\|_p^p \leq \left\| \sum_{i=1}^n A_i \right\|_p^p \leq n^{p-1} \sum_{i=1}^n \|A_i\|_p^p.$$

Note that the commutative version of Lemma 2.9 for scalars follows from the well-known Hölder inequality (see, e.g., [2, p. 88]).

The next theorem is our second main result. It can be regarded as a generalization of (1.2) in inequalities.

**Theorem 2.10.** *Let  $A_1, \dots, A_n \in C_p$  for some  $p > 0$ .*

(i) *If  $0 < p < 2$ , then*

$$\sum_{i=1}^n \left\| A_i - \frac{1}{n} \sum_{j=1}^n A_j \right\|_p^p \geq n^{\frac{p}{2}-1} \sum_{i=1}^n \|A_i\|_p^p - n \left\| \frac{1}{n} \sum_{j=1}^n A_j \right\|_p^p.$$

(ii) *If  $2 < p < \infty$ , then*

$$n^{\frac{p}{2}-1} \sum_{i=1}^n \|A_i\|_p^p - n \left\| \frac{1}{n} \sum_{j=1}^n A_j \right\|_p^p \geq \sum_{i=1}^n \left\| A_i - \frac{1}{n} \sum_{j=1}^n A_j \right\|_p^p.$$

*Proof.* Let  $0 < p < 2$ . Then

$$\begin{aligned}
& \sum_{i=1}^n \left\| A_i - \frac{1}{n} \sum_{j=1}^n A_j \right\|_p^p + n \left\| \frac{1}{n} \sum_{j=1}^n A_j \right\|_p^p \\
&= \sum_{i=1}^n \left\| \left| A_i - \frac{1}{n} \sum_{j=1}^n A_j \right|^2 \right\|_{p/2}^{p/2} + n \left\| \left| \frac{1}{n} \sum_{j=1}^n A_j \right|^2 \right\|_{p/2}^{p/2} \\
&\geq \left\| \sum_{i=1}^n \left| A_i - \frac{1}{n} \sum_{j=1}^n A_j \right|^2 + n \left| \frac{1}{n} \sum_{j=1}^n A_j \right|^2 \right\|_{p/2}^{p/2} \\
&\quad \text{(by the second inequality of Lemma 2.9(i))} \\
&= \left\| \sum_{i=1}^n |A_i|^2 \right\|_{p/2}^{p/2} \quad \text{(by equality (2.3))} \\
&\geq n^{\frac{p}{2}-1} \sum_{i=1}^n \left\| |A_i|^2 \right\|_{p/2}^{p/2} \\
&\quad \text{(by the first inequality of Lemma 2.9(i))} \\
&= n^{\frac{p}{2}-1} \sum_{i=1}^n \|A_i\|_p^p.
\end{aligned}$$

This proves the first part of the theorem. By a similar argument, one can prove the second part.  $\square$

The next theorem may be compared to Theorem 2.10. It can also be viewed as a variance of (1.2).

**Theorem 2.11.** *Let  $A_1, \dots, A_n \in C_p$  for some  $p > 0$ . Then*

$$\sum_{i=1}^n \left\| A_i - \frac{1}{n} \sum_{j=1}^n A_j \right\|_p^2 \geq n^{\frac{2}{p}-1} \sum_{i=1}^n \|A_i\|_p^2 - n \left\| \frac{1}{n} \sum_{j=1}^n A_j \right\|_p^2$$

for  $0 < p < 2$ ; and

$$n^{\frac{2}{p}-1} \sum_{i=1}^n \|A_i\|_p^2 - n \left\| \frac{1}{n} \sum_{j=1}^n A_j \right\|_p^2 \geq \sum_{i=1}^n \left\| A_i - \frac{1}{n} \sum_{j=1}^n A_j \right\|_p^2$$

for  $2 \leq p < \infty$ .



*Proof.* Let  $0 < p < 2$ . Then

$$\begin{aligned}
& \sum_{i=1}^n \left\| A_i - \frac{1}{n} \sum_{j=1}^n A_j \right\|_p^2 + n \left\| \frac{1}{n} \sum_{j=1}^n A_j \right\|_p^2 \\
&= \sum_{i=1}^n \left\| \left| A_i - \frac{1}{n} \sum_{j=1}^n A_j \right|^2 \right\|_{p/2} + n \left\| \left| \frac{1}{n} \sum_{j=1}^n A_j \right|^2 \right\|_{p/2} \\
&\leq \left\| \sum_{i=1}^n \left| A_i - \frac{1}{n} \sum_{j=1}^n A_j \right|^2 + n \left| \frac{1}{n} \sum_{j=1}^n A_j \right|^2 \right\|_{p/2} \\
&= \left\| \sum_{i=1}^n |A_i|^2 \right\|_{p/2} \quad (\text{by equality (2.3)}) \\
&\leq \left( \sum_{i=1}^n \| |A_i|^2 \|_{p/2}^{p/2} \right)^{2/p} \quad (\text{by the second inequality of Lemma 2.9(i)}) \\
&= \left( \sum_{i=1}^n \| A_i \|_p^p \right)^{2/p} \\
&\leq n^{\frac{2}{p}-1} \sum_{i=1}^n \left( \| A_i \|_p^p \right)^{2/p} \quad (\text{by a scalar version of Lemma 2.9 (ii)}) \\
&= n^{\frac{2}{p}-1} \sum_{i=1}^n \| A_i \|_p^2.
\end{aligned}$$

This proves the first part of the theorem. The second part of the theorem follows from a similar argument. The second author thanks the NSU Farquhar College of Arts and Sciences for a Mini-grant.  $\square$

**Acknowledgement.** The authors would like to express their gratitude to the referee for his/her very useful suggestions.

## REFERENCES

- [1] D. Amir, *Characterizations of Inner Product Spaces in Operator Theory: Advances and Applications*, 20, Birkhäuser Verlag, Basel, 1986.
- [2] R. Bhatia, *Matrix Analysis*, Springer-Verlag, New York, 1997.
- [3] R. Bhatia and F. Kittaneh, *Cartesian decompositions and Schatten norms*, Linear Algebra Appl. **318** (2000), 109–116.
- [4] R. Bhatia and F. Kittaneh, *Norm inequalities for partitioned operators and an application*, Math. Ann. **287** (1990), no. 4, 719–726.
- [5] S.S. Dragomir, *Some Schwarz type inequalities for sequences of operators in Hilbert spaces*, Bull. Austral. Math. Soc. **73** (2006), no. 1, 17–26.
- [6] F. Hansen and G.K. Pedersen, *Jensen's operator inequality*, Bull. London Math. Soc. **35** (2003), 553–564.

- [7] O. Hirzallah, F. Kittaneh and M.S. Moslehian, *Schatten  $p$ -norm inequalities related to a characterization of inner product spaces*, arXiv: 0801.2726v1 [math.OA].
- [8] Th.M. Rassias, *New characterizations of inner product spaces*, Bull. Sci. Math. (2) **108** (1984), no. 1, 95–99.
- [9] B. Simon, *Trace Ideals and their Applications*, Cambridge University Press, Cambridge, 1979.
- [10] F. Zhang, *On the Bohr inequality of operators*, J. Math. Anal. Appl. **333** (2007), 1264–1271.

DEPARTMENT OF MATHEMATICS, FERDOWSI UNIVERSITY OF MASHHAD, P.O. Box 1159, MASHHAD 91775, IRAN;

CENTRE OF EXCELLENCE IN ANALYSIS ON ALGEBRAIC STRUCTURES (CEAAS), FERDOWSI UNIVERSITY OF MASHHAD, IRAN.

*E-mail address:* `moslehian@ferdowsi.um.ac.ir` and `moslehian@ams.org`

FARQUHAR COLLEGE OF ARTS AND SCIENCES, NOVA SOUTHEASTERN UNIVERSITY, 3301 COLLEGE AVE, FORT LAUDERDALE, FL 33314, USA;

COLLEGE OF MATHEMATICS AND SYSTEMS SCIENCE, SHENYANG NORMAL UNIVERSITY, SHENYANG, LIAONING PROVINCE 110034, CHINA.

*E-mail address:* `zhang@nova.edu`